arXiv:1401.0407v2 [math.AP] 8 Apr 2014

CAUCHY INDEPENDENT MEASURES AND ALMOST-ADDITIVITY OF ANALYTIC CAPACITY

VLADIMIR EIDERMAN, ALEXANDER REZNIKOV, AND ALEXANDER VOLBERG

Abstract. We show that, given a family of discs centered at a chord-arc curve, the analytic capacity of a union of arbitrary subsets of these discs (one subset in each disc) is comparable with the sum of their analytic capacities. However we need that the discs in the question would be separated, and it is not clear whether the separation condition is essential or not. We apply this result to find families $\{\mu_j\}$ of measures in $\mathbb{C}$ with the following property. If the Cauchy integral operators $\mathcal{C}_{\mu_j}$ from $L^2(\mu_j)$ to itself are bounded uniformly in j, then $\mathcal{C}_\mu$, $\mu = \sum \mu_j$, is also bounded from $L^2(\mu)$ to itself.

1. Introduction

We consider two properties of families of sets and measures in the complex plane.

1.1. **Almost additivity of analytic capacity.** The *analytic capacity* $\gamma(F)$ of a compact set F in $\mathbb{C}$ is defined by the equality

$$\gamma(F) = \sup |f'(\infty)|,$$

where the supremum is taken over all analytic functions $f\colon \mathbb{C}\setminus F \to \mathbb{C}$ with $|f| \le 1$ on $\mathbb{C}\setminus F$. Here $f'(\infty) = \lim_{z\to\infty} z(f(z) - f(\infty))$. For non-compact F we set

$$\gamma(F) = \sup\{\gamma(K) : K \text{ compact}, K \subset F\}$$

[G]. For a summary of equivalent definitions the reader can see [To] and [Vo].

In the celebrated paper [To1] Tolsa established the countable semiadditivity of the analytic capacity, i. e. that

$$\gamma\big(\bigcup F_i\big) \le C \sum \gamma(F_i)$$

with an absolute constant C. But the inverse inequality does not hold in general. To see that we consider the n-th generation $E_n^{1/4}$ of the corner 1/4-Cantor set constructed in the following way. Start with the unit square (0-th generation). The j-th generation consists of 4^j squares $E_{j,k}$ with side length 4^{-j}, each square $E_{j,k}$ contains four squares of $(j+1)$-th generation, located at the corners of $E_{j,k}$, and so on. It's known [MTV] that $\gamma(\bigcup_{k=1}^{4^n} E_{n,k}) = \gamma(E_n^{1/4}) \asymp 1/\sqrt{n}$ with absolute constants of comparison; here $P \asymp Q$ means that $cP \le Q \le CP$. Positive constants c, C, a, A (possibly with indices) are not necessarily the same at each appearance. On the other hand,

$$\sum_{k=1}^{4^n} \gamma(E_{n,k}) \asymp 4^n \cdot 4^{-n} = 1.$$

Alexander Volberg was partially supported by the U.S. NSF grant DMS-0758552.

Thus, "almost additivity" $\gamma\big(\bigcup F_i\big) \asymp \sum \gamma(F_i)$ of the analytic capacity does not take place in general. N. A. Shirokov posed the question on the validity of this property for the special class of sets described in the following Theorem 1.1.

We say that Γ is a chord-arc curve, if

$$|t-s| \le A_0|z(t)-z(s)|, \quad A_0 > 1, \tag{1.1}$$

where $z(t)$ is the arc-length parametrization of Γ.

Theorem 1.1. *Let D_j be discs with centers on a chord-arc curve Γ, such that $\lambda D_j \cap \lambda D_k = \emptyset$, $j \neq k$, for some $\lambda > 1$. Let $E_j \subset D_j$ be arbitrary compact sets. Then there exists a constant $c = c(\lambda, A_0)$, such that*

$$\gamma\big(\bigcup E_j\big) \ge c \sum_j \gamma(E_j). \tag{1.2}$$

We conclude the present subsection with a useful corollary.

Corollary 1.2. *If Γ in Theorem 1.1 is the union of n chord-arc curves Γ_i with the same constant A_0, then*

$$\gamma\big(\bigcup E_j\big) \ge \frac{c}{n} \sum_j \gamma(E_j), \quad c = c(\lambda, A_0).$$

In particular, Γ might be a circle ($n = 2$). Moreover, $\{D_j\}$ might be a family of λ-separated discs (that is λD_j are disjoint), $\lambda > 1$, intersecting the same circle Γ, not necessarily with centers on Γ.

Proof. Let $D_j = D(x_j, r_j)$, and let m be such that

$$\max_{1\le k\le n} \Big\{ \sum_{j: x_j \in \Gamma_k} \gamma(E_j) \Big\} = \sum_{j: x_j \in \Gamma_m} \gamma(E_j).$$

Then by Theorem 1.1 we have

$$\gamma\big(\bigcup E_j\big) \ge \gamma\Big(\bigcup_{j: x_j \in \Gamma_m} E_j\Big) \ge c \sum_{j: x_j \in \Gamma_m} \gamma(E_j) \ge \frac{c}{n} \sum_j \gamma(E_j).$$

For the last assertion it is sufficient to prove that if all discs D_j intersect a semicircle T, then there is another chord-arc curve $\widetilde{\Gamma}$ with a constant $A_0 = A_0(\lambda)$ containing all centers of D_j. We may assume that there are more than one disc D_j. Fix j, and let a_j, b_j be the points of intersection of T and the circle $\partial(\lambda' D_j)$, where $\lambda' = \frac{1+\lambda}{2}$ (for two discs D_j it might be only one such a point). We replace the arc of T with the endpoints a_j, b_j by two line segments $x_j a_j$ and $x_j b_j$ (and by the only one line segment in the case of one point of intersection). We claim that the obtained curve is chord-arc. Indeed, since $|a_j - b_j| \ge c(\lambda) r_j$, the condition (1.1) holds with $A_0 = A_0(\lambda)$ for any points $z(t), z(s) \in \lambda' D_j$, and for any two points in $\widetilde{\Gamma}$ situated on T. If $z(t) \in \lambda' D_i$, $z(s) \in \lambda' D_j$, $i \neq j$, then both parts of (1.1) are comparable with $|x_i - x_j|$. To demonstrate this assertion we notice that the inequality $|z_1 - z_2| > (\lambda - \lambda')(r_i + r_j)$, $z_1 \in \lambda' D_i$, $z_2 \in \lambda' D_j$, implies the relations $|z_1 - z_2| \asymp |x_i - x_j|$ and

$$t - s \le C(r_i + |a_i - b_j| + r_j) \le C' |x_i - x_j|.$$

Similar arguments yield (1.1) when one point is situated on T. □

Open question. It is not clear if the theorem is true or not when $\lambda = 1$.

1.2. **Cauchy independence of families of measures.** We use the results in the previous subsection to investigate the property of measures described below.

We call a finite Borel measure with compact support in the complex plane *a Cauchy operator measure* if the Cauchy operator $\mathcal{C}_\mu$ is bounded from $L^2(\mu)$ to itself wth norm at most 1.

The first natural question is how to interpret the "definition" of $\mathcal{C}_\mu$ as

$$\mathcal{C}_\mu f(z) = \int \frac{f(\xi)d\mu(\xi)}{\xi - z}.$$

One of the ways is to consider the so-called ε-truncations, defined by

$$\mathcal{C}^\varepsilon_\mu f(z) = \int_{\varepsilon<|\xi-z|<\varepsilon^{-1}} \frac{f(\xi)d\mu(\xi)}{\xi - z}.$$

We now say that $\mathcal{C}_\mu$ is bounded as an operator from $L^2(\mu)$ to itself if the ε-truncations are bounded from $L^2(\mu)$ to itself uniformly in ε. Moreover, by the norm of $\mathcal{C}_\mu$ we understand the $\sup_{\varepsilon>0} \|\mathcal{C}^\varepsilon_\mu\|_\mu =: \|\mathcal{C}_\mu\|_\mu$, where $\|\mathcal{C}^\varepsilon_\mu\|_\mu$ is the norm of $\mathcal{C}^\varepsilon_\mu$ as an operator from $L^2(\mu)$ to itself. We encourage the reader to look for other interpretations in [NTrV1], [To] and [Vo].

The following important fact (which we will repeatedly use) demonstrates the connection between the analytic capacity and boundedness of the Cauchy operator [To1, To2, To, Vo]: for every compact set F in $\mathbb{C}$,

$$\gamma(F) \asymp \sup\{\|\mu\| : \operatorname{supp}\mu \subset F,\ \mu \in \Sigma,\ \|\mathcal{C}_\mu\|_\mu \le 1\}, \tag{1.3}$$

where Σ is the class of nonnegative Borel measures μ such that $\mu(D(x,r)) \le r$ for every disc $D(x,r) := \{z \in \mathbb{C} : |z-x| < r\}$.

We call a collection $\{\mu_j\}$ of finite positive Borel measures with compact supports *C-Cauchy independent measures* if a) $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \le 1$ (Cauchy operator measures) and b) $\|\mathcal{C}_\mu\|_\mu \le C < \infty$ for $\mu = \Sigma_j \mu_j$. We will call such collection *Cauchy independent* if it is C-Cauchy independent for some finite C.

The family $\{\mu_j\}$ can be finite or infinite. Two Cauchy operator measures are always Cauchy independent with an absolute constant C. A short proof of this nontrivial fact is given in [NToV, Proposition 3.1]. So, a finite family is always Cauchy independent for a sufficiently large constant C. But our main interest is in situations, when infinite families are independent (or when C is independent of the number of measures). The main result is the following

Theorem 1.3. *Suppose that $\lambda > 1$, and measures μ_j are supported on compact sets E_j lying in discs D_j such that λD_j are disjoint. We also assume that measures μ_j are extremal in the following sense: $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \le 1$ and $\|\mu_j\| \asymp \gamma(E_j)$ with absolute comparison constants. Let $\mu = \sum_j \mu_j$ and $E = \cup E_j$. Then this family is Cauchy independent if and only if for any disc B,*

$$\mu(B) \le C_0 \gamma(B \cap E)\,. \tag{1.4}$$

Remarks. 1. In Section 3 we prove that the condition (1.4) with any disc B is necessary for the bound $\|\mathcal{C}_\mu\|_\mu \le C$ without any additional assumptions on the structure of μ. Example 5.1 given in Section 5 shows that this condition alone is not sufficient even if μ consists of countably many pieces μ_j, and each of μ_j gives a bounded Cauchy operator with a uniform bound. Thus, additional conditions on the structure of μ are needed. The example of such assumptions on μ which seems

reasonable is given in Theorem 1.3, where supports of μ_j are located in separated discs.

2. In general one cannot discard the requirement that the measures μ_j in Theorem 1.3 are extremal – see Example 5.2 in Section 5.

3. On pp. 125–129, 135–146 of the paper of Tolsa [To1] it is proved that under the conditions of Theorem 1.3, there exists a piece of measure μ, namely $\mu' := \chi_{E'} \cdot \mu$, $E' \subset E$, such that $\mu'(E) \geq c\,\mu(E)$, and $\|\mathcal{C}_{\mu'}\|_{\mu'} \leq C < \infty$, where $c > 0$ and C are constants depending only on parameters in Theorem 1.3. This fact is far from being trivial, it is used in [To1] to approach Painlevé's conjecture. In other words, to prove that under the assumptions of Theorem 1.3 a "good portion" of μ generates a bounded Cauchy operator is a highly non-trivial problem. It is remarkable that the whole measure μ has, in fact, such a property.

As a corollary of Theorem 1.3 we derive the following independence theorem.

Theorem 1.4. *Let μ_j be measures supported on compacts E_j lying in discs D_j such that λD_j are disjoint ($\lambda > 1$), and let $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \leq 1$. If for any disc B,*

$$\sum_j \gamma(B \cap E_j) \leq C_1 \gamma(B \cap E), \quad E = \bigcup E_j, \tag{1.5}$$

then the norm $\|\mathcal{C}_\mu\|_\mu$ is bounded, and the bound depends only on a comparison constant C_1 and on λ. Here, as above, $\mu = \Sigma_j \mu_j$.

Note that unlike Theorem 1.3, this result does not need the additional condition that measures μ_j are extremal. On the other hand, (1.5) is not necessary for the boundedness of $\mathcal{C}_\mu$. For example, if μ_j are just 2-dimensional Lebesgue measures on the squares E_j, $j = 1, 2, \ldots$, defined below in Example 5.1, the operator $\mathcal{C}_\mu$ is obviously bounded, but (1.5) does not hold.

Unlike Theorem 1.1, Theorems 1.3, 1.4 do not have any assumptions on the location of discs D_j. However, condition (1.5) is not completely independent of a geometry of discs. Theorem 1.1 states that if λ-separated discs are situated along a chord-arc curve, then the almost additivity of the analytic capacity takes place. We are going to prove a statement which is converse in the following sense: almost additivity of analytic capacity in the form of inequality (1.5) together with certain additional assumptions imply that our discs have to "line-up" along a good (Ahlfors-David regular) curve.

By $\mathcal{H}^1$ we denote the 1-dimensional Hausdorff measure. A set $G \subset \mathbb{C}$ is called Ahlfors-David (AD) regular if

$$c\,r \leq \mathcal{H}^1(G \cap B(x, r)) \leq Cr, \quad x \in G,\ 0 < r \leq \operatorname{diam} G,$$

with some positive constants c, C.

Corollary 1.5. *Suppose that λ-separated ($\lambda > 1$) discs $D_j = D(x_j, r_j)$ and subsets $E_j \subset D_j$ are such that (a) (1.5) holds, (b) $\gamma(E_j) \asymp r_j$, (c) the set $\mathcal{T} := \bigcup_j \partial D_j$ is AD regular. Then there exists an AD regular curve which intersects all discs D_j.*

Without any of assumptions (a) – (c) Corollary 1.5 is incorrect – see Proposition 5.3.

To prove Theorems 1.3, 1.4, we will need only the special case of Theorem 1.1 when all discs D_j intersect a real line or a circle, that is the last assertion of Corollary 1.2. In this case there is a short proof based only on some classical facts in complex

analysis. We give this proof in the next Section 2. Theorem 1.3 is proved in Section 3, and Theorem 1.4 with Corollary 1.5 in Section 4. In Section 5 we give the examples mentioned above. Section 6 contains the proof of Theorem 1.1 in the full generality, which is completely different from the proof in Section 2. The main tool of this proof is Melnikov–Menger's curvature of a measure. All necessary definitions are given in Section 6. In the last Section 7 we formulate an open question.

2. Almost-additivity of analytic capacity: string of beads attached to the real line

A result close to the theorem below for some special sets $\{E_j\}_{j=1}^{\infty}$ was proved (but not stated) in [NV]. Here we use the approach via the Marcinkiewicz function, the approach in [NV] was a bit more complicated. Unlike [NV], we do not need any special size properties of these sets.

Theorem 2.1. *Let D_j be discs such that $\lambda D_j \cap \lambda D_k = \emptyset$, $j \neq k$, for some $\lambda > 1$, and each disc D_j has a non-empty intersection with the real line $\mathbb{R}$. Let $E_j \subset D_j$ be arbitrary compact sets. Then there exists a constant $c = c(\lambda) > 0$, such that*

$$\gamma\big(\bigcup E_j\big) \geq c \sum_j \gamma(E_j).$$

Proof. It is enough to prove the result for finite families of indices j. We first notice that $\gamma_j := \gamma(E_j) \leq \gamma(D_j) = r_j$, where r_j is the radius of D_j. Let y_j be the center of the chord $\mathbb{R} \cap D_j$. Denote $\lambda' := \frac{1+\lambda}{2}$. For each j we draw a horizontal line segment $L_j \subset \lambda' D_j$ with center at y_j and with the analytic capacity $b(\lambda)\gamma_j$, where $b(\lambda) = \frac{\sqrt{\lambda'^2-1}}{4}$. Thus, the length ℓ_j of L_j satisfies $\ell_j = 4b(\lambda)\gamma_j \leq r_j\sqrt{\lambda'^2-1}$. In particular, the whole segment L_j lies in $\lambda' D_j$.

Next, let f_j be the function that gives the capacity of E_j. Also let φ_j be the function that gives the capacity of L_j. Then we have

$$\varphi_j(z) = \int_{L_j} \frac{\varphi_j(x)}{x-z} dx, \quad \int_{L_j} \varphi_j(x)dx = b(\lambda)\gamma_j.$$

Positive functions $\varphi_j(x)$ have a uniform bound $\|\varphi_j\|_\infty \leq A$ with an absolute constant A. In particular, if $\mathcal{F}$ is any subset of indices j we have

$$\left|\mathrm{Im} \sum_{j\in\mathcal{F}} \varphi_j(z)\right| \leq A \int_{\bigcup_{j\in\mathcal{F}} L_j} \frac{|\mathrm{Im} z|}{|t-z|^2}\, dt \leq \pi A\,, \quad \forall z \in \mathbb{C}\,. \tag{2.1}$$

Remark. It is important here that the intervals L_j are situated on the real line (or at least are not far away from $\mathbb{R}$). For any $M > 0$ one can easily construct a chord-arc curve and discs centered on it such that the left hand side in (2.1) exceeds M. This is the obstacle for extension of these arguments to chord-arc curves.

Our next goal is to find a family $\mathcal{F}$ of indices and absolute positive constants a_1, a_2, such that the following two assertions hold:

$$\sum_{j\in\mathcal{F}} \gamma_j \geq a_1 \sum_j \gamma_j\,, \tag{2.2}$$

$$\sum_{j\in\mathcal{F}} |f_j(z) - b(\lambda)^{-1}\varphi_j(z)| \leq a_2\,, \quad \forall z \in \mathbb{C} \setminus \bigcup_{j\in\mathcal{F}} (E_j \cup L_j). \tag{2.3}$$

Let us finish the proof of the theorem, taken these assertions as granted (for a short while). Let $F := \sum_{j\in\mathcal{F}} f_j$. Combining (2.1) and (2.3) we get $|\operatorname{Im} F(z)| \le C_1(\lambda)$, $z \in \mathbb{C}\setminus(\bigcup_{j\in\mathcal{F}} E_j)$. Hence, the function $F_1 := e^{iF} - 1$ is bounded in $\mathbb{C}\setminus(\bigcup_{j\in\mathcal{F}} E_j)$ by a constant $C(\lambda)$. Since $F(\infty) = 0$, we have $|F_1'(\infty)| = |F'(\infty)| = \sum_{j\in\mathcal{F}} \gamma_j$. Thus,

$$\gamma\Big(\bigcup_{j\in\mathcal{F}} E_j\Big) \ge \frac{1}{C(\lambda)} \sum_{j\in\mathcal{F}} \gamma_j\,.$$

Combine this with (2.2). We obtain, that

$$\gamma\Big(\bigcup_{j} E_j\Big) \ge \gamma\Big(\bigcup_{j\in\mathcal{F}} E_j\Big) \ge a_3 \sum_{j} \gamma_j\,,$$

and Theorem 1.1 would be proved. So we are left to chose the family $\mathcal{F}$ such that (2.2), (2.3) hold.

By the Schwartz lemma in the form we borrow from [G, p. 12–13], we have

$$|f_j(z) - b(\lambda)^{-1}\varphi_j(z)| \le \frac{A r_j \gamma_j}{\operatorname{dist}(z, E_j\cup L_j)^2}\,, \quad z\notin E_j\cup L_j\,. \tag{2.4}$$

Denote

$$Q_i := \lambda' D_i\,, \quad g_i := \sum_{j:\, j\ne i} \frac{r_j\gamma_j}{D(Q_j,Q_i)^2}\,,$$

where $D(Q_i,Q_j) := \operatorname{dist}(Q_i,Q_j)$.

Remark. We do not need this, but for the sake of explanation, let us define a function $g = \sum g_j \chi_{Q_j\cap\mathbb{R}}$. This function is often called a Marcinkiewicz function. The main trick with Marcinkiewicz functions is to integrate them with respect to a suitable measure. What in fact happens next is that we integrate it with respect to Lebesgue measure on $\mathbb{R}$.

The important point is that we can estimate $\sum_i g_i\gamma_i$. In fact,

$$\begin{aligned}\sum_i g_i\gamma_i &= \sum_i \gamma_i \sum_{j:\, j\ne i} \frac{r_j\gamma_j}{D(Q_j,Q_i)^2} = \sum_j r_j\gamma_j \sum_{i:\, i\ne j} \frac{\gamma_i}{D(Q_i,Q_j)^2}\\ &\le \sum_j r_j\gamma_j \sum_{i:\, i\ne j} \frac{r_i}{D(Q_i,Q_j)^2} \le A_0 \sum_j r_j\gamma_j r_j^{-1} = A_0 \sum_j \gamma_j\,.\end{aligned}$$

In the last estimate we used that

$$\sum_{i:\, i\ne j} \frac{r_i}{D(Q_i,Q_j)^2} \le A_1 \int_{t:|t-y_j|\ge r_j} \frac{1}{|t-y_j|^2}\,dt \le \frac{2A_1}{r_j}\,, \quad A_1 = A_1(\lambda),$$

since the length of $\mathbb{R}\cap Q_j$ exceeds $c(\lambda)r_j$, and $D(Q_i,Q_j) \ge c'(\lambda)|t-y_j|$, $t\in\mathbb{R}\cap Q_i$.

Now we apply the Tchebychev inequality. Denote $I^* := \{i:\ g_i > 10A_0\}$, $I_* := \{i:\ g_i \le 10A_0\}$. We immediately see that

$$\sum_{j\in I_*} \gamma_j \ge \frac{9}{10} \sum_j \gamma_j\,. \tag{2.5}$$

Obviously, by (2.4) for every index i we have

$$\sum_{j:\, j\ne i} |f_j(z) - b(\lambda)^{-1}\varphi_j(z)| \le A g_i\,, \quad z\in Q_i\,.$$

This estimate and the choice of I_* imply that

$$\sum_{j:\, j\neq i,\ j\in I_*} |f_j(z) - b(\lambda)^{-1}\varphi_j(z)| \le C(\lambda) g_i \le 10A_0 A\,, \quad \forall i \in I_*\,,\ \forall z \in Q_i\,.$$

But all functions $|f_i|, |\varphi_i|$ are bounded by 1 in $\mathbb{C} \setminus (E_i \cup L_i)$. Therefore, the last inequality implies the estimate

$$\sum_{j:\, j\in I_*} |f_j(z) - b(\lambda)^{-1}\varphi_j(z)| \le 10A_0 A + b(\lambda)^{-1} + 1 =: a_2 \qquad \forall z \in Q_i \setminus (E_i \cup L_i)\,,\ \forall i \in I_*. \tag{2.6}$$

The function $\sum_{j\in I_*}(f_j - b(\lambda)^{-1}\varphi_j)$ is analytic in $\mathbb{C} \setminus \bigcup_{i\in I_*}(E_i \cup L_i)$ and vanishes at infinity. Therefore, (2.6) implies (2.3) if we put $\mathcal{F} := I_*$. Assertion (2.2) is proved in (2.5), and the proof of Theorem 2.1 is completed. □

Corollary 2.2. *The statement of Theorem 2.1 remains true if discs D_j intersect a circle instead of the real line.*

Proof. It is sufficient to consider the unit circle. There are at most K discs D_j with radii $r_j \ge 1/30$, intersecting the unit circle Γ, where K is an absolute constant. Hence, we may assume that $r_j < 1/30$. Moreover, as in Corollary 1.2, we may restrict ourself by discs intersecting the left semicircle T. Let $h(z) := i\frac{1+z}{1-z}$ be the conformal mapping of Γ onto the real line. Let E be a compact subset of $G^0 := \{z : \operatorname{dist}(z, T) < 1/10\}$, and $\mathcal{E} := h(E)$. We prove that

$$\gamma(E) \asymp \gamma(\mathcal{E}) \tag{2.7}$$

with absolute constants of comparison. Certainly, this relation is a consequence of the general result by Tolsa [To3] about stability of the analytic capacity under bilipschitz maps. But we prefer a direct elementary proof (which possibly is not new) without Tolsa's very difficult result. Pick any function $f(z)$ such that $f(z)$ is holomorphic and $|f(z)| < 1$ in $\mathbb{C} \setminus E$. Define the sets

$$G := \{z : \operatorname{dist}(z, T) < 1/5\}, \quad \mathcal{G} := h(G), \quad \mathcal{G}^0 := h(G^0),$$

and the function $F(w) := f(g(w))g'(w)$, where $g(w) = \frac{w-i}{w+i}$ is the inverse of h. Clearly, $|F(w)| \le C_1$ as $w \in \mathcal{G} \setminus \mathcal{E}$, and the length of $\partial\mathcal{G}$ does not exceed C_2, where C_1, C_2 are absolute constants. Fix $w \in \mathcal{G}^0 \setminus \mathcal{E}$, and let L^0 be a closed curve in $\mathcal{G}^0 \setminus \mathcal{E}$ which encloses $\mathcal{E}$ but not w, and is oriented in such a way that w is "on the left". If $\partial\mathcal{G}$ is oriented in the same way, by the Cauchy formula we have

$$F(w) = \frac{1}{2\pi i}\int_{\partial\mathcal{G}} \frac{F(\xi)}{\xi - w}\, d\xi + \frac{1}{2\pi i}\int_{L^0} \frac{F(\xi)}{\xi - w}\, d\xi =: F_1(w) + F_2(w).$$

Since $|\xi - w|$ exceeds an absolute constant as $\xi \in \partial\mathcal{G}$, $|F_1(w)| < C$ as $w \in \mathcal{G}^0 \setminus \mathcal{E}$. But F is bounded in this domain as well. Hence, $F_2(w)$ is bounded in $\mathcal{G}^0 \setminus \mathcal{E}$ and holomorphic in $\mathbb{C}\setminus\mathcal{E}$. By the maximum principle, $F_2(w)$ is holomorphic and bounded by an absolute constant in $\mathbb{C} \setminus \mathcal{E}$. Let L be a contour in $\mathcal{G} \setminus \mathcal{E}$ inclosing $\mathcal{E}$. Since $F_1(w)$ is holomorphic in $\mathcal{G}$, we have $\int_L F_1(w)\, dw = 0$. Hence,

$$\int_L F_2(w)\, dw = \int_L F(w)\, dw = \int_L f(g(w))\, dg(w) = \int_{g(L)} f(z)\, dz.$$

Thus, $\gamma(E) \le C\gamma(\mathcal{E})$. The similar arguments yield the inverse inequality, if we start with the compact $\mathcal{E}$ (we may assume that $\mathcal{E} \subset \mathcal{G}^0$). Therefore, (2.7) is proved. Corollary 2.2 is a direct consequence of (2.7) and Theorem 2.1. □

3. Proof of Theorem 1.3

3.1. **Necessity of the condition** (1.4)**.** Suppose that $\|\mathcal{C}_\mu\|_\mu \le C < \infty$ and $\operatorname{supp}\mu \subset E$. One can easily see that $\|\mathcal{C}_{\mu|B}\|_{\mu|B} \le C < \infty$ for any disc B. Moreover, boundedness of $\mathcal{C}_\mu$ implies that $\alpha\mu \in \Sigma$ with α depending only on C, see for example [Da]. Thus, the measure $c\mu|B$, $c = c(C) > 0$, participates in the right hand side of (1.3) with $F = B \cap E$, and we get (1.4).

3.2. **Sufficiency of the condition** (1.4)**.** The following result was proved in [NToV], although was not formulated explicitly (see the last three pages of Section 3 in [NToV]).

Theorem 3.1. *Suppose that $\{D_j\}$ are discs on the plane and the dilated discs λD_j, $\lambda > 1$, are disjoint. Let ν, σ be two positive measures supported in $\bigcup_j D_j$ such that $c_1\nu(D_j) \le \sigma(D_j) \le c_2\nu(D_j)$, $0 < c_1 < c_2 < \infty$. Suppose also that the Cauchy operators $\mathcal{C}_{\sigma_j}$, $\sigma_j = \sigma|_{D_j}$, are uniformly bounded. Then if ν is a Cauchy operator measure, then $\alpha\sigma$ is also a Cauchy operator measure with a constant α depending only on c_1, c_2, and λ.*

We need some preliminary constructions and notations. Here is an easy lemma.

Lemma 3.2. *For any circle T and any disc B,*

$$\gamma(T \cap B) \asymp \mathcal{H}^1(T \cap B)$$

with absolute constants of comparison.

Now we define new L_j. We need a number $N = N(\lambda)$, which is defined as follows. Recall that $\lambda > 1$ and $\lambda' = \frac{1+\lambda}{2}$. Let a disc D with radius r be given. We place the circle of radius $A_\lambda r$, $A_\lambda := \min(1, \lambda' - 1)/1000$, in the center of D, and N circles that touch $\partial(\lambda' D_j)$ from within, of the same radius as the first circle, and on equal distance from each other. We also require that circles do not intersect. By L we denote the union of all circles. Let N be a minimal integer such that the following holds.

If a disc B intersects D and $\mathbb{C} \setminus \lambda D$, then at least one circle from L lies inside B.

Clearly, such $N = N(\lambda) < \infty$ exists, and this property remains valid if we reduce the radii of circles in L. Since $\gamma(\text{circle}) \asymp \mathcal{H}^1(\text{circle})$, we have the obvious lemma:

Lemma 3.3. *For the set L defined above, $\gamma(L) \asymp \mathcal{H}^1(L)$, where the comparison constants can depend only on N.*

Let L_j be the union of circles associated with D_j. We have chosen the number of circles in each L_j, but we have a freedom to choose their size. We redefine the radius so that

$$\mathcal{H}^1(L_j) = A_\lambda(N+1)\gamma(E_j). \tag{3.1}$$

Then, in particular, $\mathcal{H}^1(\text{one circle}) = \frac{1}{N+1}\mathcal{H}^1(L_j) = A_\lambda\gamma(E_j) \le A_\lambda r_j$, since $\gamma(E_j) \le \gamma(D_j) = r_j$.

We need the following lemma.

Lemma 3.4. *Fix an index j. Let B be a disc such that at least one circle from L_j lies inside B. Then $\gamma(L_j) \asymp \gamma(L_j \cap B)$ with constants depending only on λ. In particular this is true if $D_j \subset B$, or if B intersects D_j and $\mathbb{C} \setminus \lambda D_j$.*

Proof. Indeed, by semiadditivity of γ we have $\gamma(L_j) \leq A(N+1)\gamma(\text{central circle}) \leq A(N+1)\gamma(L_j \cap B)$. □

Lemma 3.5. *For any disc B the following relation holds with constants depending only on λ:*

$$\gamma\Big(\bigcup_{j:D_j\subset B} L_j\Big) \asymp \gamma\Big(\bigcup_{j:D_j\subset B} L_j \cap B\Big).$$

Proof. By semiadditivity of γ,

$$\gamma\Big(\bigcup_{D_j\subset B} L_j\Big) \leq A\bigg(\gamma\Big(\bigcup_{\lambda' D_j\subset B} L_j\Big) + \gamma\Big(\bigcup_{D_j\subset B,\ \lambda' D_j\not\subset B} L_j\Big)\bigg).$$

The first term is the same as $\gamma(\bigcup_{\lambda' D_j\subset B} L_j \cap B)$. For the second, we notice that $L_j \cap B \subset \lambda' D_j$. Since discs $\frac{\lambda}{\lambda'}\lambda' D_j$ are pairwise disjoint, we may apply Corollary 2.2 with just a new dilation constant $\lambda_{new} := \frac{\lambda}{\lambda'}$. Thus,

$$\begin{aligned}\gamma\Big(\bigcup_{D_j\subset B,\ \lambda' D_j\not\subset B} L_j \cap B\Big) &\geq c \sum_{D_j\subset B,\ \lambda' D_j\not\subset B} \gamma(L_j \cap B)\\ &\geq c_1 \sum_{D_j\subset B,\ \lambda' D_j\not\subset B} \gamma(L_j) \geq c_2\gamma\Big(\bigcup_{D_j\subset B,\ \lambda' D_j\not\subset B} L_j\Big),\end{aligned}$$

which finishes the proof. The second inequality uses Lemma 3.4. □

For a given disc B denote by $\mathcal{J} = \mathcal{J}(B)$ the set of indices $\mathcal{J} := \{j : D_j \cap B \neq \emptyset \text{ and } D_j \not\subset B\}$.

Lemma 3.6. *Suppose that a disc B intersects more than one D_j. Then*

$$\gamma\Big(\bigcup_{\mathcal{J}} L_j\Big) \asymp \gamma\Big(\bigcup_{\mathcal{J}} L_j \cap B\Big).$$

with comparison constants depending only on λ.

Proof. Here again we will use Corollary 2.2 of Theorem 2.1. Since B intersects more than one D_j, it cannot be contained in λD_j, $j \in \mathcal{J}$. Thus, it contains at least one circle from L_j for each $j \in \mathcal{J}$ (it follows from the choice of N). Call this circle C_j. We apply Corollary 2.2 to D_j with dilation constant λ to get the estimate

$$\gamma\Big(\bigcup_{\mathcal{J}} L_j \cap B\Big) \geq c\sum_{\mathcal{J}} \gamma(L_j \cap B) \geq \sum_{\mathcal{J}} \gamma(C_j) \geq c_1 \sum_{\mathcal{J}} \gamma(L_j) \geq c_2\gamma\Big(\bigcup_{\mathcal{J}} L_j\Big),$$

which finishes the proof. □

Finally, we need the following notation. Fix a disc B. Denote

$$F_j = \begin{cases} E_j, & D_j \subset B \\ \emptyset, & D_j \not\subset B. \end{cases}, \quad F = \bigcup F_j.$$

Remark. A disc B will be free to change in what follows. The constants in further inequalities will never depend on B.

Our next goal is to prove that under assumptions of Theorem 1.3, the inequality

$$\gamma\big(\bigcup L_j \cap B\big) \geq c \sum \gamma(L_j \cap B)$$

holds with a universal constant c (universality means that c will not depend on the disc B). We need the following two lemmas.

We fix a small positive absolute constant ε. The choice of smallness will be clear from what follows.

Lemma 3.7 (The first case)**.** *Suppose that $\gamma(F) \leq \varepsilon\gamma(E \cap B)$. Then there exists a constant c, that can depend only on N, ε, and other universal constants, such that*

$$\gamma\big(\bigcup L_j \cap B\big) \geq c \sum \gamma(L_j \cap B).$$

Proof. Suppose that B intersects only one λD_j. Then the $\bigcup$ and the $\sum$ have only one term, and there is nothing to prove. So, we can assume that B intersects at least two of λD_j's. Notice also that by this assumption, by the fact that λD_i are pairwise disjoint, and by the choice of N, if B intersects D_j then at least one circle from L_j lies inside B. Let $\mathcal{J}$ be as in Lemma 3.6. Using Lemma 3.4 and Corollary 2.2 we get

$$\sum_{\mathcal{J}} \gamma(L_j) \leq A_1 \sum_{\mathcal{J}} \gamma(L_j \cap B) \leq A_2 \gamma\big(\bigcup_{\mathcal{J}} L_j \cap B\big)\,. \tag{3.2}$$

On the other hand, by the assumption of Theorem 1.3,

$$\sum_{D_j \subset B} \gamma(L_j) \leq C \sum_{D_j \subset B} \gamma(E_j) \leq C' \sum_{D_j \subset B} \mu_j(D_j) \leq C'\mu(B) \leq C'C_0\gamma(E \cap B). \tag{3.3}$$

Also with an absolute constant A,

$$\gamma(E \cap B) \leq A\Big(\gamma(F) + \gamma\big(\bigcup_{\mathcal{J}} E_j \cap B\big)\Big) \leq \varepsilon A\gamma(E \cap B) + A\gamma\big(\bigcup_{\mathcal{J}} E_j \cap B)\big).$$

Thus, if ε is small enough (notice that the smallness depends only on A), we have

$$\gamma(E \cap B) \leq C\gamma\big(\bigcup_{\mathcal{J}} E_j \cap B)\big). \tag{3.4}$$

Therefore, combining (3.3), (3.4), and (3.2), we obtain

$$\begin{aligned} \sum_{D_j \subset B} \gamma(L_j) &\leq C\gamma\big(\bigcup_{\mathcal{J}} E_j \cap B)\big) \leq C_1 \sum_{\mathcal{J}} \gamma(E_j \cap B) \\ &\leq C_2 \sum_{\mathcal{J}} \gamma(L_j) \leq C_3 \gamma\big(\bigcup_{\mathcal{J}} L_j \cap B\big). \end{aligned} \tag{3.5}$$

Now combine (3.2) and (3.5) to get

$$\gamma\big(\bigcup L_j \cap B\big) \geq \gamma\big(\bigcup_{\mathcal{J}} L_j \cap B\big) \geq c \sum_{D_j \subset B} \gamma(L_j) + c \sum_{\mathcal{J}} \gamma(L_j) = c \sum_{D_j \cap B \neq \emptyset} \gamma(L_j)\,. \tag{3.6}$$

Obviously,

$$\gamma\big(\bigcup L_j \cap B\big) \geq \gamma\big(\bigcup_{\mathcal{J}_1} L_j \cap B\big), \quad \mathcal{J}_1 := \{j : D_j \cap B = \emptyset,\ L_j \cap B \neq \emptyset\}. \tag{3.7}$$

For $j \in \mathcal{J}_1$ we again consider the new dilation constants $\lambda_{new} := \frac{\lambda}{\lambda'}$, and discs $D'_j := \lambda' D_j$. The discs $\lambda_{new} D'_j$, $j \in \mathcal{J}_1$, are disjoint, and D'_j intersects B for $j \in \mathcal{J}_1$. By Corollary 2.2 applied to $L_j \cap B$, $j \in \mathcal{J}_1$, playing the roles of E_j, we get

$$\gamma\big(\bigcup_{\mathcal{J}_1} L_j \cap B\big) \geq c \sum_{\mathcal{J}_1} \gamma(L_j \cap B)\,. \tag{3.8}$$

The combination of (3.6)–(3.8) finishes the proof. □

Lemma 3.8 (The second case)**.** *Suppose that $\gamma(F) \geq \varepsilon\gamma(E \cap B)$ with ε from the previous lemma. Then there exists a universal constant c such that*

$$\gamma\big(\bigcup L_j \cap B\big) \geq c \sum \gamma(L_j \cap B).$$

Proof. By Theorem 2.1 or rather Corollary 2.2 we need only to prove the inequality

$$\gamma\big(\bigcup_{D_j \subset B} L_j \cap B\big) \geq c \sum_{D_j \subset B} \gamma(L_j \cap B). \tag{3.9}$$

Using the assumption of our lemma as well as the conditions (1.4) and $\|\mu_j\| \asymp \gamma(E_j)$ of Theorem 1.3, we get

$$\gamma(F) \geq \varepsilon\gamma(E \cap B) \geq \varepsilon c \mu(B) \geq \varepsilon c \sum_{D_j \subset B} \mu_j(B) \geq \varepsilon c' \sum_j \gamma(F_j). \tag{3.10}$$

By ν we denote the measure on F participating in (1.3) for which $\|\nu\| \asymp \gamma(F)$. Denote $d\nu_j = \chi_{F_j} d\nu$. Then $\mathcal{C}_{\nu_j}$ is bounded on $L^2(\nu_j)$ (with norm at most 1), and (1.3) yields the estimate

$$\|\nu_j\| \leq C\gamma(F_j) \leq C_1\gamma(L_j) \leq C_2\mathcal{H}^1(L_j) =: C_2\ell_j.$$

We call j *good* if $D_j \subset B$ and $\|\nu_j\| \geq \tau\ell_j$. The choice of τ will be clear from the next steps. However, we want to emphasize now that this choice will be universal. By (3.10) we have:

$$\begin{aligned}\varepsilon c' A^{-1} \sum \gamma(F_j) &\leq A^{-1}\gamma(F) \leq \|\nu\| = \sum \|\nu_j\| \\ &\leq C_2 \sum_{j \text{ is good}} \ell_j + \tau \sum_{D_j \subset B} \ell_j \leq C_2 \sum_{j \text{ is good}} \ell_j + C_3\tau \sum \gamma(F_j)\end{aligned}$$

(in the last inequality we use (3.1)). Therefore,

$$\sum_{j \text{ is good},\ D_j \subset B} \ell_j \geq c \sum \gamma(F_j) \geq c_1 \sum_{D_j \subset B} \ell_j. \tag{3.11}$$

Actually, τ is chosen exactly here. We see that it indeed depends only on universal constants such as A and ε. Recall that C_j denotes the central circle of each L_j. We set

$$d\sigma_g := \sum_{j \text{ is good},\ D_j \subset B} \chi_{C_j} d\mathcal{H}^1, \quad d\nu_g := \sum_{j \text{ is good},\ D_j \subset B} d\nu_j\,.$$

Then for good j, $\sigma_g(D_j) = \mathcal{H}^1(C_j) \asymp \mathcal{H}^1(L_j) = \ell_j \asymp \nu_g(D_j)$. In the last relation the comparison constants can depend on previous universal constants and τ. Operators $\mathcal{C}_{\sigma_g|D_j}$ are uniformly bounded. By the choice of ν, the operator $\mathcal{C}_{\nu_g}$ is bounded as well with norm at most 1. Thus, we may apply Theorem 3.1 and conclude that

$\mathcal{C}_{\sigma_g}$ is also bounded with a certain absolute bound of the norm. Therefore, using (3.11), we get

$$\gamma(\bigcup_{D_j\subset B} L_j) \geq \gamma(\bigcup_{j \text{ is good, } D_j\subset B} L_j) \geq c\|\sigma_g\| \geq c_1 \sum_{j \text{ is good, } D_j\subset B} \ell_j \geq c_2 \sum_{D_j\subset B} \ell_j.$$

In Lemma 3.5 we have proved that

$$\gamma(\bigcup_{D_j\subset B} L_j) \asymp \gamma(\bigcup_{D_j\subset B} L_j \cap B).$$

Moreover, for every j such that $D_j \subset B$, we have

$$\ell_j = \mathcal{H}^1(L_j) \asymp \mathcal{H}^1(L_j \cap B) \asymp \gamma(L_j \cap B).$$

Thus, we obtain (3.9), and Lemma 3.8 is proved. □

The main Theorem of [NV] says:

Theorem 3.9. *Let $L \subset \mathbb{R}^2$, be a compact set of positive and finite Hausdorff measure $\mathcal{H}^1$, and let $\sigma = \mathcal{H}^1|L$. Then $\mathcal{C}_\sigma$ is bounded if and only if there exists a finite constant C_0 such that $\sigma(B \cap L) \leq C_0\gamma(B \cap L)$ for any disc B.*

Starting with the main assumption of Theorem 1.4 (the inequality $\mu(B) \leq C_0\gamma(B\cap E)$ for any disc B) we proved in Lemmas 3.7, 3.8 that the uniform in B almost-additivity of γ holds for the union of all sets $\{L_j \cap B\}$. Namely, we proved that the following holds for any B with uniform positive c_2:

$$\gamma(B \cap L) \geq c_1 \sum_j \gamma(B \cap L_j) \geq c_2 \sum_j \sigma(B \cap L_j) = c_2\sigma(B \cap L)\,, \tag{3.12}$$

where $\sigma := \mathcal{H}^1|L$. Hence the measure σ satisfies Theorem 3.9. So the boundedness of Cauchy operator on the union of circles is obtained. The measures $\sigma|L_j$ and μ_j are supported on $\lambda' D_j$, the discs $\frac{\lambda}{\lambda'} \cdot (\lambda' D_j)$ are disjoint, and $\sigma(L_j) \asymp \mu_j$ (see (3.1)). We may apply Theorem 3.1 again to establish the boundedness of $\mathcal{C}_\mu$ in $L^2(\mu)$.

4. Proof of Theorem 1.4 and Corollary 1.5

Proof of Theorem 1.4. We would like to reduce the assumptions of Theorem 1.4 to conditions of Theorem 1.3. For every E_j, choose an "extremal" measure μ'_j, supported on E_j and such that $\|\mathcal{C}_{\mu'_j}\|_{\mu'_j} \leq 1$, and $c'_1\mu'_j(E_j) \leq \gamma(E_j) \leq c'_2\mu'_j(E_j)$, $0 < c'_1 < c'_2 < \infty$ (the existence of such measures μ'_j follows from (1.3)).

Now let us consider new measures

$$\widetilde{\mu}_j := \mu_j + \mu'_j\,, \quad \widetilde{\mu} = \sum_j \widetilde{\mu}_j\,.$$

We prove that these new measures satisfy all assumptions of Theorem 1.3 (up to an absolute constant). Indeed, we saw in Section 3.1 that the assumption $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \leq 1$ implies the inequality $\mu_j(E_j) \leq C\gamma(E_j)$. Hence, the new measures $\widetilde{\mu}_j$ also satisfy the "extremality" condition (μ_j did not satisfy it in general):

$$c_1\widetilde{\mu}_j(E_j) \leq \gamma(E_j) \leq c_2\widetilde{\mu}_j(E_j), \quad 0 < c_1 < c_2 < \infty.$$

Moreover, since two Cauchy operator measures are always Cauchy independent (see Section 1.2), $\|\mathcal{C}_{\widetilde{\mu}_j}\|_{\widetilde{\mu}_j} \le C$. Now the relation (1.3) implies that $\widetilde{\mu}(B \cap E_j) = \widetilde{\mu}_j(B \cap E_j) \le C\gamma(B \cap E_j)$ for every disc B. Therefore, by (1.5) we have

$$\widetilde{\mu}(B) = \sum \widetilde{\mu}(B \cap E_j) \le C \sum \gamma(B \cap E_j) \le C_0 \gamma(B \cap E), \quad C_0 = C_0(C_1).$$

where the latter is the condition (1.4) of Theorem 1.3 for $\widetilde{\mu}$.

We apply Theorem 1.3 to $c\widetilde{\mu}_j$ and $c\widetilde{\mu} = c\sum_j \widetilde{\mu}_j$ with a sufficiently small absolute constant $c > 0$, and get that $\|\mathcal{C}_{\widetilde{\mu}}\|_{\widetilde{\mu}} \le C(C_1, \lambda)$. But $\mu = \sum_j \mu_j$ is a part of μ, and hence $\|\mathcal{C}_\mu\|_\mu \le C(C_1, \lambda)$. □

Proof of Corollary 1.5. Recall the reader that above we built L_j for each D_j, and each L_j contains a certain "central circle". Let $\mathcal{T}'$ be the union over j of central circles in L_j. By (3.1), the radii of these circles are comparable with r_j. Since $\mathcal{T}$ is AD regular and discs D_j are λ-separated, $\mathcal{T}'$ is AD regular as well (with another constants c, C). We have proved in the previous section that the Cauchy operator $\mathcal{C}_\sigma$, $\sigma := \mathcal{H}^1|L$, is bounded from $L^2(\sigma)$ to itself. Hence, the operator $\mathcal{C}_{\sigma'}$, $\sigma' := \mathcal{H}^1|\mathcal{T}'$, is bounded too. By the theorem of Mattila, Melnikov and Verdera [MMV], the set $\mathcal{T}'$ is contained in an AD regular curve. □

5. Examples

We saw in Section 3 that the condition

$$\mu(B) \le C_0 \gamma(B \cap E) \text{ for every disc } B \tag{5.1}$$

is necessary for the boundedness of the Cauchy operator $\mathcal{C}_\mu$ with any Borel measure μ. It is not difficult to see that this condition alone is not enough for the boundedness of $\mathcal{C}_\mu$. Indeed, let $\mu_n^{1/4}$ be the probability measure uniformly distributed on the set $E_n^{1/4}$ defined in Introduction. Let $\mu^{1/4}$ be the weak limit of some weakly convergent subsequence $\{\mu_{n_k}^{1/4}\}$, $E^{1/4} = \bigcap E_n^{1/4}$, E is the initial unit square, and $\mu := \mu^{1/4} + \mathcal{H}^2|E$. Then μ satisfies (5.1), but $\mathcal{C}_\mu$ is unbounded – see for example [MT, MTV]. We are going to demonstrate more: in general the condition (5.1) is not sufficient for the boundedness even if μ consists of countably many pieces, and each of them gives a bounded Cauchy operator.

Example 5.1. *There exists a family of measures $\{\mu_j\}_{j=0}^\infty$, supported on squares E_j, with the following properties: (a) $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \le 1$; (b) $\|\mu_j\| \asymp \gamma(E_j)$; (c) $2E_j \cap 2E_k = \emptyset$, $j \ne k$, $j, k \ge 1$; (d) the measure $\mu = \sum_{j=0}^\infty \mu_j$ satisfies (5.1); (e) $\|\mathcal{C}_\mu\|_\mu = \infty$.*

Proof. We use the idea of David-Semmes (see [VE, Example 8.7] for more detailed exposition). Let $N_0 = 0$, and let $\{N_k\}_{k=0}^\infty$ be a sequence of natural numbers such that $N_{k+1} - N_k \to \infty$ as $k \to \infty$. Start the construction with the unit square E_0 and make $N_1 - N_0$ steps of the construction of the corner 1/4-Cantor set $E^{1/4}$. We get $4^{N_1 - N_0}$ squares with side length 4^{-N_1}. Choose one (any) of them, denote it by Q_1, and continue the construction only with this square. Other $4^{N_1-N_0} - 1$ squares are the sets E_j which already have been defined. For the chosen square Q_1 we make next $N_2 - N_1$ steps of the construction of $E^{1/4}$, obtaining $4^{N_2-N_1}$ squares with side length 4^{-N_2}. Again, continue the construction only for one of them, say, for a square Q_2, and so on.

Let μ_j, $j = 0, 1, \dots$, be the 2-dimensional measure uniformly distributed on E_j such that $\|\mu_j\| = c\ell_j$, where ℓ_j is the side length of E_j, and the absolute constant

c is chosen in such a way that $\|\mathcal{C}_{\mu_j}\|_{\mu_j} = 1$. Then properties (a), (b), (c) are obvious. To demonstrate (d) we notice that $E := \bigcup_{j\geq 0} E_j$ is equal to E_0, and thus $\gamma(B\cap E) \asymp \operatorname{diam}(B\cap E) =: d_0$. On the other hand, for any $j \geq 0$ and $d_j := \operatorname{diam}(B\cap E_j)$, we have $\mu(B\cap E_j) \leq c\ell_j^{-1} d_j^2 < C d_j$ (the density of μ_j is equal to c/ℓ_j). Hence, $\mu(B\cap E) < C\sum_{j=0}^{\infty} d_j \asymp d_0$, and (d) is established.

Finally, to prove (e), we apply the operator $\mathcal{C}_\mu$ to the characteristic functions χ_{Q_k}, $k = 0, 1, \dots$ We have

$$\|\mathcal{C}_\mu(\chi_{Q_k})\|_{L^2(\mu)} = \|\mathcal{C}_{\mu|Q_k}(\mathbf{1})\|_{L^2(\mu)} \geq \|\mathcal{C}_{\mu|Q_k}(\mathbf{1})\|_{L^2(\mu|Q_k)}.$$

But

$$\|\mathcal{C}_{\mu|Q_k}(\mathbf{1})\|^2_{L^2(\mu|Q_k)} \geq c(N_{k+1} - N_k)4^{-N_k}$$

with an absolute constant c – see [MT]. Hence, $\|\mathcal{C}_\mu\|_\mu \geq c(N_{k+1} - N_k) \to \infty$, and (e) is proved. □

Remark that the measures $\{\mu_j\}_{j=1}^{\infty}$ satisfy all assumptions of Theorem 1.3 except (5.1). Therefore, we have to add μ_0 and change the structure of μ.

Now we demonstrate that the condition $\|\mu_j\| \asymp \gamma(E_j)$ in Theorem 1.3 is essential.

Example 5.2. *There exists a family of measures $\{\mu_j\}_{j=1}^{\infty}$ with the following properties: (a) $\|\mathcal{C}_{\mu_j}\|_{\mu_j} \leq 1$; (b) $\|\mu_j\| \leq C\gamma(E_j)$, where $E_j = \operatorname{supp}\mu_j$, and E_j is either a square or a disc; (c) $2E_j \cap 2E_k = \emptyset$, $j \neq k$; (d) the measure $\mu = \sum_{j=1}^{\infty}\mu_j$ satisfies (5.1); (e) $\|\mathcal{C}_\mu\|_\mu = \infty$.*

Proof. We use the same construction as in Example 5.1, but with the following modifications. 1. The initial square E_0 now is not included into the collection $\{E_j\}$ of squares; thus, the squares are separated. 2. Besides the same squares E_j and measures μ_j, as in Example 5.1, we add additional discs $\widetilde{E}_j$ and measures $\widetilde{\mu}_j$ to satisfy (5.1) (otherwise (5.1) does not hold after exclusion E_0).

As before, set $N_0 = 0$, choose a sequence $\{N_k\}_{k=0}^{\infty}$ of natural numbers such that $N_{k+1} - N_k \to \infty$ as $k \to \infty$, and make $N_1 - N_0$ steps of the construction of the corner 1/4-Cantor set $E^{1/4}$ starting with the unit square E_0. Let $E_{n,k}$, $k = 0, \dots, 4^n$, be the squares forming the nth generation in this construction (not all of them will be included into $\{E_j\}$). In each square $E_{n,k}$, $n = 0, \dots, N_1$, $k = 1, \dots, 4^n$, place the disc $\widetilde{D}_{n,k}$ concentric with $E_{n,k}$, and with radius $\ell_n/10 = 4^{-n}/10$. On $\widetilde{D}_{n,k}$ we uniformly distribute a measure $\mu_{n,k}$ with $\|\mu_{n,k}\| = 2^{-n}\cdot 4^{-n}$. Then for n bigger than a certain absolute n_0 we automatically have $\|\mathcal{C}_{\mu_{n,k}}\|_{\mu_{n,k}} \leq 1$. For $n \leq n_0$ we might need a small absolute positive c' such that $\|\mathcal{C}_{c'\mu_{n,k}}\|_{c'\mu_{n,k}} \leq 1$. We put then $\widetilde{\mu}_{n,k} := c'\mu_{n,k}$ and achieve that always $\|\mathcal{C}_{\widetilde{\mu}_{n,k}}\|_{\widetilde{\mu}_{n,k}} \leq 1$. After $N_1 - N_0$ steps choose one (any) of the squares $E_{N_1,k}$, and denote it by Q_1. Other $4^{N_1-N_0} - 1$ squares $E_{N_1,k}$ and discs $\widetilde{D}_{n,k}$, $n = 0, \dots, N_1$, $k = 1, \dots, 4^n$, are the sets E_j which already have been defined. As before, $\mu_{N_1,k}$ is the 2-dimensional measure uniformly distributed on $E_{N_1,k}$ such that $\|\mu_{N_1,k}\| = c\ell_{N_1}$ and $\|\mathcal{C}_{\mu_{N_1,k}}\|_{\mu_{N_1,k}} = 1$.

For the chosen square Q_1 we continue the construction and make the next $N_2 - N_1$ steps of the construction of $E^{1/4}$, obtaining the squares $E_{N_2,k}$, $k = 1, \dots, 4^{N_2-N_1}$, with side length $\ell_{N_2} := 4^{-N_2}$ and with measures $\mu_{N_2,k}$ such that $\|\mu_{N_2,k}\| = c\ell_{N_2}$. Besides these squares, we get discs $\widetilde{D}_{n,k}$, $n = N_1, \dots, N_2$, $k = 1, \dots, 4^{n-N_1}$ of radii $4^{-n}/10$, concentric with $E_{n,k}$ and supporting the measures $\widetilde{\mu}_{n,k}$, $\|\widetilde{\mu}_{n,k}\| = c'2^{-n}\cdot 4^{-n}$. Again, continue the construction only for one of them, and so on.

We have to prove only (d). Fix a disc B. Suppose that $\widetilde{D}_{0,1} \subset B$. Since $\mu(B) \le \|\mu\| \le C$ and $\gamma(B \cap E) \ge \gamma(\widetilde{D}_{0,1}) = 1/10$, (5.1) holds. Suppose now that $\widetilde{D}_{0,1} \not\subset B$, and that B contains at least one disc $\widetilde{D}_{n,k}$. Choose a maximal disc in B, say, $\widetilde{D}_{n_1,k_1}$. Then "the parent" $E_{n_1-1,k_1'}$ of the square E_{n_1,k_1} (that is the square of the previous generation containing E_{n_1,k_1}) does not lie in B: otherwise $\widetilde{D}_{n_1,k_1}$ would not be a maximal disc in B. Set $G_1 := E_{n_1-1,k_1'} \cap B$. Now choose a maximal disc $\widetilde{D}_{n_2,k_2} \subset (B \setminus G_1)$ (if such a disc exist). Its "parent" $E_{n_2-1,k_2'}$ and the set G_1 are disjoint. Hence, $E_{n_2-1,k_2'} \not\subset B$. Set $G_2 := E_{n_2-1,k_2'} \cap B$. Continuing in such a way, we obtain a sequence of sets G_j with the following properties. (i) $2G_i \cap 2G_j = \emptyset$, $i \ne j$, and one may place them into λ-separated discs, $\lambda > 1$. (ii) For each G_j,

$$\mu(G_j) \le C\ell_{n_j} = C4^{-n_j} \asymp \gamma(\widetilde{D}_{n_j,k_j}) \le \gamma(G_j).$$

(iii) All squares $E_{N_i,k}$ and discs $\widetilde{D}_{n,k}$ contained in B are contained in $\cup_j G_j$.

Also, it might be some discs $\widetilde{D}_{n,k}$ and squares $E_{N_i,k}$ intersecting B. Each of them form a separate set $G_j := \widetilde{D}_{n,k} \cap B$ or $E_{N_i,k} \cap B$. These sets are λ-separated as well, and

$$\mu(G_j) \le C \operatorname{diam}(G_j) \le C'\gamma(G_j)$$

(see the proof of Example 5.1). Moreover, all sets G_j satisfy the conditions of Corollary 2.2 which yields the estimate

$$\mu(B) \le \sum_j \mu(G_j) \le C \sum_j \gamma(G_j) \le C'\gamma(B \cap E),$$

and the proof is completed. □

Now we demonstrate the sharpness of Corollary 1.5.

Proposition 5.3. *Without any of assumptions (a)–(c) Corollary 1.5 is incorrect.*

Proof. **1.** Suppose that the condition (a) in Corollary 1.5 is missed. Then we may use the same sets $\{E_j\}_{j=1}^{\infty}$ as in Example 5.1, only without the initial square E_0. As D_j we take discs containing E_j and slightly bigger than E_j. A proof of (b) and (c) is not difficult, and we leave it to the reader. At the same time,

$$\text{(length of any curve connecting all discs in } Q_k)/4^{-N_k} \to \infty \text{ as } k \to \infty \tag{5.2}$$

(the squares Q_k are defined in the proof of Example 5.1).

2. Consider the case when (b) is missed. Let D_j be the discs $\widetilde{D}_{n,k}$ in Example 5.2, enumerated in the order of non-increase of their radii r_j. In each D_j we place the disc E_j concentric with D_j and of radius $\frac{1}{20}4^{-j}$. To prove (a), fix a disc B. For discs E_j intersecting ∂B, (1.5) holds by Corollary 2.2 (with $B \cap E_j$ as E_j). Let $j_0 := \{\min j : E_j \subset B\}$. Then

$$\begin{aligned}\sum_{j:B_j \subset B} \gamma(B \cap E_j) &\le \sum_{j=j_0}^{\infty} \gamma(E_j) = \sum_{j=j_0}^{\infty} \frac{1}{20}4^{-j} \\ &= \frac{1}{15}4^{-j_0} = \frac{8}{3}\gamma(E_{j_0}) \le \frac{8}{3}\gamma(B \cap E),\end{aligned}$$

and (a) is proved. The proof of (c) is easy and we skip it. At the same time, (5.2) holds for discs D_j in Q_k.

3. Finally, suppose that (c) is missed. The counterexample in this case is not based on Cantor-type constructions. Given $\ell > 0$, $N \in \mathbb{N}$, $N \geq 4$, consider the square $Q_\ell = [0, \ell] \times [0, \ell]$ and points

$$x_i = \frac{\ell}{N-1} i, \quad y_j = \frac{\ell}{N-1} j, \quad i, j = 0, 1, \dots, N-1.$$

Let E_{ij} be discs centered at the points (x_i, y_j) with radii ℓ/N^2, and let $\mathcal{E} := \bigcup_{i,j} E_{ij}$, $D_{ij} := 2E_{ij}$. Fix a disc B. Set $\mathcal{E}_B = \{\bigcup E_{ij} : E_{ij} \subset B\}$ ($\mathcal{E}_B$ might be empty). If $\mathcal{E}_B \neq \emptyset$, we have

$$\left| \int \frac{d(\mathcal{H}^1|\partial\mathcal{E}_B)(\xi)}{\xi - z} \right| < C, \quad z \in \mathbb{C}.$$

Hence,

$$\gamma(\mathcal{E} \cap B) \geq \gamma(\mathcal{E}_B) \geq c\mathcal{H}^1(\partial\mathcal{E}_B) = c' \sum_{E_{ij} \subset B} \gamma(E_{ij}) \geq c'' \sum \gamma(E_{ij} \cap B).$$

If $\mathcal{E}_B = \emptyset$, (1.5) holds as well (for example, by Corollary 2.2). At the same time, the length of any curve, intersecting all discs D_{ij}, is at least $C\ell N$.

Now we construct a series of squares Q_{ℓ_k} with $\ell_k = \frac{1}{10} 2^{-k}$, $N_k = k\, 2^k$, centered at points $1/k^2$, and the corresponding sets $\mathcal{E}^{(k)}$ and discs $D_{ij}^{(k)}$, $E_{ij}^{(k)}$. One may place $\mathcal{E}^{(k)}$ into λ-separated discs centered at points $1/k^2$. Set $E := \bigcup_k \mathcal{E}^{(k)}$ (that is E is the union of all discs $E_{ij}^{(k)}$). By Theorem 2.1,

$$\gamma(B \cap E) \geq c \sum_k \gamma(B \cap \mathcal{E}^{(k)}).$$

Thus, to prove (1.5), we have to establish almost additivity of γ for each $\mathcal{E}^{(k)}$ separately, that was done above. The validity of (b) is obvious. But all discs $D_{ij}^{(k)}$ in E cannot be connected by an AD regular curve. □

6. Proof of Theorem 1.1

It is known that a compact chord-arc curve is a bi-lipschitz image of a straight segment, see [Po], Chapter 7. On the other hand analytic capacity can be only finitely distorted by bi-lipschitz maps. This is a difficult result by X. Tolsa, [To3]. So if we allow the separation constant $\lambda > 1$ to depend on the Lipschitz constant of our chord-arc curve (so, the separation of the discs to be large if the constant of the curve is large), then we can obtain Theorem 1.1 directly from Theorem 2.1. However, we want to avoid the dependence of the separation constant on the chord-arc constant. Then we need another proof, which follows.

The Melnikov–Menger curvature of a positive Borel measure μ in $\mathbb{C}$ is defined as

$$c^2(\mu) = \iiint \frac{1}{R^2(x, y, z)}\, d\mu(x)\, d\mu(y)\, d\mu(z),$$

where $R(x, y, z)$ is the radius of the circle passing through points $x, y, z \in \mathbb{C}$, with $R(x, y, z) = \infty$ if x, y, z lie on the same straight line (in particular, if two of these points coincide). This notion was introduced by Melnikov [M]. The following relation characterizes the analytic capacity in terms of the curvature of a measure [To1], [To2, p. 104], [Vo], [To] : for any compact set F in $\mathbb{C}$,

$$\gamma(F) \asymp \sup\{\mu(F) :\ \operatorname{supp}\mu \subset F,\ \mu \in \Sigma,\ c^2(\mu) \leq \mu(F)\}, \tag{6.1}$$

where Σ is the class of measures of linear growth defined in (1.3).

Lemma 6.1 (Main Lemma)**.** *Let $D_j = D(x_j, r_j)$ be discs with centers on a chord-arc curve Γ, such that $\lambda D_j \cap \lambda D_k = \emptyset$, $j \neq k$, for some $\lambda > 1$. Let μ_j be positive measures with the following properties: (1) $\operatorname{supp}\mu_j \subset D_j$; (2) $\mu_j(B_j) =: \|\mu_j\| \leq r_j$. Then for $\mu = \sum \mu_j$ we have*

$$c^2(\mu) \leq \sum_j c^2(\mu_j) + C\|\mu\|, \quad C = C(\lambda, A_0), \tag{6.2}$$

where A_0 is the constant of Γ.

At the beginning let us show that Theorem 1.1 is a direct consequence of Main Lemma and (6.1).

Proof of Theorem 1.1. Consider measures μ_j participating in (6.1) for $F = E_j$, $j = 1, \dots$ Then $\mu(D(x,r)) \leq Cr$ for any disc D, where $C = C(A_0)$ and $\mu = \sum \mu_j$. To prove this assertion, we fix a disc $D = D(x,r)$ and divide all discs D_j onto two groups: $\mathcal{D}_1 := \{D_j : D_j \cap D \neq \emptyset,\ r_j \leq r\}$, $\mathcal{D}_2 := \{D_j : D_j \cap D \neq \emptyset,\ r_j > r\}$. Since Γ is chord-arc, $\sum_{D_j \in \mathcal{D}_1} r_i \leq Cr$, $C = C(A_0)$. It is easy to see that $\#\mathcal{D}_2 \leq 6$. Hence,

$$\mu(D) \leq \sum_{D_j \in \mathcal{D}_1} \mu(D_j) + \sum_{D_j \in \mathcal{D}_2} \mu(D_j \cap D) \leq \sum_{D_j \in \mathcal{D}_1} r_j + 6\mu(D) < Cr.$$

Furthermore, Main Lemma implies the inequality $c^2(\mu) \leq C\|\mu\|$, $C = C(\lambda, A_0)$. Thus, the measure $c\mu$ with an appropriate constant c depending on λ, A_0, participates in (6.1) for $F = E = \cup E_j$. So, $\gamma(E) \geq c\|\mu\|$, that implies Theorem 1.1. $\square$

Proof of Lemma 6.1. It is enough to consider the case of a finite set of discs B_j, $j = 1, \dots, N$. We assume that these discs are enumerated in the order of increase of the natural parameters of their centers.

Let Γ_j be arcs of Γ such that $\Gamma_j \subset D_j$ and $\mathcal{H}^1(\Gamma_j) = \mu(D_j)$. Let $\sigma_j := \mathcal{H}^1|\Gamma_j$ and $\sigma := \sum \sigma_j$, so that $\sigma(D_j) = \mu(D_j)$. Obviously,

$$c^2(\mu) = \Big(\sum_j \iiint_{D_j^3} + \iiint_{\mathbb{C}^3 \setminus \bigcup_j D_j^3} \Big) \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) =: I_1 + I_2.$$

Since $I_1 = \sum_j c^2(\mu_j)$, we have to estimate only I_2. Our proof is based on the comparison of I_2 and the corresponding integral with respect to σ:

$$\bar{I}_2 := \iiint_{\mathbb{C}^3 \setminus \bigcup_j D_j^3} \frac{1}{R^2(x,y,z)}\, d\sigma(x)\, d\sigma(y)\, d\sigma(z).$$

Notice that

$$\bar{I}_2 < c^2(\sigma) \leq C\|\sigma\|, \quad C = C(A_0). \tag{6.3}$$

The last inequality is a consequence of two well-known facts. (a) The boundedness of the Cauchy operator $\mathcal{C}_{\mathcal{H}^1|\Gamma}$ on chord-arc curves – see [MV, p. 330]. In particular,

$$\|\mathcal{C}_\sigma^\varepsilon \mathbf{1}\|^2_{L^2(\sigma)} \leq \|\mathcal{C}_{\mathcal{H}^1|\Gamma} \chi_{\cup \Gamma_j}\|^2_{L^2(\mathcal{H}^1|\Gamma)} \leq C\|\chi_{\cup \Gamma_j}\|^2_{L^2(\mathcal{H}^1|\Gamma)} = C\|\sigma\|, \quad \varepsilon > 0,$$

where C depends only on A_0. (b) The connection between the curvature of a measure and the norm of a Cauchy potential:

$$\|\mathcal{C}_\mu^\varepsilon \mathbf{1}\|^2_{L^2(\mu)} = \frac{1}{6} c_\varepsilon^2(\mu) + O(\|\mu\|)$$

for any measure $\mu \in \Sigma$ uniformly in ε – see for example [To2]. Here $c^2_\varepsilon(\mu)$ is the truncated version of $c^2(\mu)$ defined in the same way as $c^2_\varepsilon(\mu)$, but the triple integral is taken over the set $\{(x,y,z) \in \mathbb{C}^3 : |x-y|, |y-z|, |x-z| > \varepsilon\}$. This equality with $\mu = c\,\sigma \in \Sigma$, and the previous relations imply (6.3).

Obviously,

$$I_2 = \Big(\iiint_{\Omega_1} + \iiint_{\Omega_2} \Big) \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) =: I_{2,1} + I_{2,2}\,,$$

where

$$\Omega_1 := \{D_j \times D_k \times D_l : j = k \neq l \vee j \neq k = l \vee j = l \neq k\},$$
$$\Omega_2 := \{D_j \times D_k \times D_l : j \neq k,\ k \neq l,\ j \neq l\}.$$

To estimate the integral over Ω_1, it's sufficient to consider the subset

$$\Omega_1' := \{D_j \times D_k \times D_l : j \neq k = l\}.$$

For $x \in D_j = D(x_j, r_j),\ y, z \in D_k,\ j \neq k$, we have

$$2R(x,y,z) \geq |x-y| \geq c(r_j + r_{j+1} + \cdots + r_k), \quad c = c(\lambda, A_0)$$

(here we assume that $j < k$; the case $k < j$ is analogous). Then

$$\iiint_{\Omega_1'} \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) \leq C \Big[\sum_{j=1}^{N-1} \|\mu_j\| \sum_{k=j+1}^{N} \frac{\|\mu_k\|^2}{(r_j + r_{j+1} + \cdots + r_k)^2}$$
$$+ \sum_{j=1}^{N-1} \|\mu_{N+1-j}\| \sum_{k=j+1}^{N} \frac{\|\mu_{N+1-k}\|^2}{(r_{N+1-j} + r_{j+1} + \cdots + r_{N+1-k})^2} =: C[S_{N,1} + S_{N,2}].$$

Estimates for both terms on the right are the same. We estimate $S_{N,1}$ using the induction with respect to N.

1. $N = 2$. Then

$$S_{N,1} = \|\mu_1\| \cdot \frac{\|\mu_2\|^2}{(r_1 + r_2)^2} \leq \|\mu_1\| \leq \|\mu_1\| + \|\mu_2\|.$$

2. Suppose that the inequality

$$S_{N,1} = \sum_{j=1}^{N-1} \|\mu_j\| \sum_{k=j+1}^{N} \frac{\|\mu_k\|^2}{(r_j + \cdots + r_k)^2} \leq \|\mu_1\| + \cdots + \|\mu_N\| \tag{6.4}$$

holds for some $N \geq 2$. For $N+1$ discs we have

$$S_{N+1,1} = S_{N,1} + \sum_{j=1}^{N} \|\mu_j\| \frac{\|\mu_{N+1}\|^2}{(r_j + \cdots + r_{N+1})^2}$$
$$\leq S_{N,1} + \|\mu_{N+1}\|^2 \sum_{j=1}^{N} \frac{r_j}{(r_j + \cdots + r_{N+1})^2}.$$

The last sum is dominated by the integral

$$\int_0^\infty \frac{dt}{(r_{N+1} + t)^2} = \frac{1}{r_{N+1}}.$$

Hence,

$$S_{N+1,1} \leq S_{N,1} + \|\mu_{N+1}\|^2 / r_{N+1} \leq \|\mu_1\| + \cdots + \|\mu_{N+1}\|.$$

Thus, we proved (6.4) and therefore estimated the triple integral over Ω_1.

By symmetry,

$$\iiint_{\Omega_2} \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) = 6 \iiint_{\Omega_2'} \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z),$$

where $\Omega_2' := \{D_j \times D_k \times D_l : j < k < l\}$. Moreover, we may restrict ourself by the integration over

$$\Omega_{2,1}' := \{D_j \times D_k \times D_l : j < k < l,\ r_j + \cdots + r_k \ge \tfrac{1}{2}(r_j + \cdots + r_l)\}.$$

Indeed, if we prove the inequality

$$\iiint_{\Omega_{2,1}'} \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) \le C\|\mu\| \tag{6.5}$$

with $C = C(\lambda, A_0)$, then using the inverse parametrization of Γ, we get the same estimate for the triple integral over

$$\Omega_{2,2}' := \{D_j \times D_k \times D_l : j < k < l,\ r_k + \cdots + r_l \ge \tfrac{1}{2}(r_j + \cdots + r_l)\}$$

(here we use the same numeration of discs as before). Since $\iiint_{\Omega_2'} \le \iiint_{\Omega_{2,1}'} + \iiint_{\Omega_{2,2}'}$, (6.4) and (6.5) imply (6.2).

Fix indices j, k, l. For any triples $(x,y,z), (x',y',z') \in D_j \times D_k \times D_l$, the sine of the angle between the intervals (y,z) and (y',z') does not exceed

$$C\,\frac{r_k + r_l}{r_k + \cdots + r_l}, \quad C = C(\lambda, A_0).$$

For the angle between the intervals (x,z) and (x',z') we have $C\,\frac{r_j + r_l}{r_j + \cdots + r_l}$. Denote by α, α' the angles at z, z' of the triangles x,y,z and x',y',z', correspondingly. Since $\sin(\alpha+\beta+\gamma) \le \sin\alpha + \sin\beta + \sin\gamma$ as $\alpha, \beta, \gamma \in [0,\pi]$, we get the estimate

$$\sin\alpha < \sin\alpha' + C\,\frac{r_k + r_l}{r_k + \cdots + r_l} + C\,\frac{r_j + r_l}{r_j + \cdots + r_l}.$$

Hence,

$$\frac{1}{R(x,y,z)} = \frac{2\sin\alpha}{|x-y|} < \frac{C}{|x'-y'|}\left[2\sin\alpha' + \frac{r_k + r_l}{r_k + \cdots + r_l} + \frac{r_j + r_l}{r_j + \cdots + r_l}\right].$$

Therefore,

$$\begin{aligned}\iiint_{\Omega_2} \frac{1}{R^2(x,y,z)}\, d\mu(x)\, d\mu(y)\, d\mu(z) \le C\Bigg[&\iiint_{\Omega_2'} \frac{1}{R^2(x',y',z')}\, d\sigma(x')\, d\sigma(y')\, d\sigma(z') \\ &+ \sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \sum_{k=j+1}^{l-1} \frac{r_k^2\, \|\mu_k\|\, \|\mu_l\|}{(r_j + \cdots + r_k)^2 (r_k + \cdots + r_l)^2} \\ &+ \sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \sum_{k=j+1}^{l-1} \frac{r_l^2\, \|\mu_k\|\, \|\mu_l\|}{(r_j + \cdots + r_k)^2 (r_k + \cdots + r_l)^2} \\ &+ \sum_{j=1}^{N-2} \|\mu_j\| r_j^2 \sum_{l=j+2}^{N} \sum_{k=j+1}^{l-1} \frac{\|\mu_k\|\, \|\mu_l\|}{(r_j + \cdots + r_k)^2 (r_j + \cdots + r_l)^2}\Bigg] =: C[I + S^{(1)} + S^{(2)} + S^{(3)}].\end{aligned}$$

By (6.3), $I \le c^2(\sigma) \le C\|\sigma\|$. We estimate each of sums separately. Write $S^{(1)}$ as

$$S^{(1)} = \sum_{j=1}^{N-2} \|\mu_j\| \sum_{k=j+1}^{N-1} \sum_{l=k+1}^{N} \frac{r_k^2 \, \|\mu_k\| \, \|\mu_l\|}{(r_j + \cdots + r_k)^2 (r_k + \cdots + r_l)^2} \, .$$

Since the inner sum over l does not exceed

$$\frac{r_k^2 \, \|\mu_k\|}{(r_j + \cdots + r_k)^2} \int_{r_k}^{\infty} \frac{dx}{x^2} = \frac{r_k \, \|\mu_k\|}{(r_j + \cdots + r_k)^2},$$

we get the estimate

$$\begin{aligned} S^{(1)} &\le \sum_{j=1}^{N-2} \|\mu_j\| \sum_{k=j+1}^{N-1} \frac{r_k \, \|\mu_k\|}{(r_j + \cdots + r_k)^2} \le \sum_{k=2}^{N-1} \sum_{j=1}^{k-1} \frac{r_j \, r_k \, \|\mu_k\|}{(r_j + \cdots + r_k)^2} \\ &\le \sum_{k=2}^{N-1} \|\mu_k\| \sum_{j=1}^{k-1} r_k \int_{r_k}^{\infty} \frac{dx}{x^2} = \sum_{k=2}^{N-1} \|\mu_k\| < \|\mu\| \, . \end{aligned} \tag{6.6}$$

Now we will use the possibility to consider only those k for which $r_j + \cdots + r_k \ge \frac{1}{2}(r_j + \cdots + r_l)$ (the set of such k can be empty). Suppose that the last inequality holds for $p \le k \le l-1$. Then

$$\begin{aligned} &\sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \sum_{k=p}^{l-1} \frac{r_l^2 \, \|\mu_k\| \, \|\mu_l\|}{(r_j + \cdots + r_k)^2 (r_k + \cdots + r_l)^2} \\ &\qquad \le 4 \sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \sum_{k=p}^{l-1} \frac{r_l^2 \, \|\mu_k\| \, \|\mu_l\|}{(r_j + \cdots + r_l)^2 (r_k + \cdots + r_l)^2} \\ &\qquad\qquad \le 4 \sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \frac{r_l \, \|\mu_l\|}{(r_j + \cdots + r_l)^2} \end{aligned}$$

(we estimate the sum with respect to k in the same way as above). We may deal with the last double sum as in (6.6), or notice that this sum does not exceed

$$\sum_{j=1}^{N-2} \|\mu_j\| \left[1 + \sum_{l=j+1}^{N-1} \frac{r_l \, \|\mu_l\|}{(r_j + \cdots + r_l)^2} \right],$$

and use (6.6) directly. Finally,

$$S^{(3)} \le \sum_{j=1}^{N-2} \|\mu_j\| \sum_{l=j+2}^{N} \frac{r_j^2 \, \|\mu_l\|}{(r_j + \cdots + r_l)^2 r_j} < \sum_{j=1}^{N-2} \|\mu_j\| < \|\mu\|.$$

Lemma 6.1 is proved. □

7. Question on super-additivity

We make more accurate the question posed in Section 1. In Theorems 1.1, 2.1 discs were λ-separated, where $\lambda > 1$. But what if they are just disjoint? Namely, let D_j be circles with centers on a chord-arc curve with constant A_0 (or even on the

real line $\mathbb{R}$), such that $D_j \cap D_k = \emptyset$, $j \neq k$. Let $E_j \subset D_j$ be arbitrary compact sets. Is it true that there exists a constant $c = c(\lambda, A_0) > 0$, such that

$$\gamma(\bigcup E_j) \geq c \sum_j \gamma(E_j)\,?$$

We cannot either prove or construct a counter-example to this claim.

References

[Da] G. David, *Wavelets and singular integrals on curves and surfaces,* Lecture Notes in Math. **1465**, Springer-Verlag, Berlin, 1991.

[G] J. Garnett, Analytic Capacity and Measure. Springer-Verlag. 1972.

[MT] J. Mateu, X. Tolsa, *Riesz transforms and harmonic Lip_1-capacity in Cantor sets*, *Proc. London Math. Soc.* **89** (2004), no. 3, 676–696.

[MTV] J. Mateu, X. Tolsa, J. Verdera, *The planar Cantor sets of zero analytic capacity and the local $T(b)$-theorem*, J. Amer. Math. Soc. **16** (2003), no. 1, 19–28.

[MMV] P. Mattila, M.S. Melnikov, J. Verdera, *The Cauchy integral, analytic capacity, and uniform rectifiability,* Ann. of Math. (2) **144** (1996), 127–136.

[M] M. Melnikov, *Analytic capacity: discrete approach and curvature of the measure*, Mat. Sb. **186** (1995), 827–846.

[MV] M. Melnikov, J. Verdera, *A geometric proof of the L^2 boundedness of the Cauchy integral on Lipschitz graphs*, Internat. Math. Res. Notices 1995, no. 7, 325–331.

[NV] F. Nazarov, A. Volberg, *Analytic capacity of the portion of continuum and a question of T. Murai*, Proc. of Symposia in Pure Math, v. 79 (2008), pp. 279–292. The volume dedicated to Vladimir Maz'ya's 70th anniversary.

[NTrV1] F. Nazarov, S. Treil and A. Volberg, *Weak type estimates and Cotlar inequalities for Calderón-Zygmund operators in nonhomogeneous spaces,* Int. Math. Res. Notices **9** (1998), 463–487.

[NToV] F. Nazarov, X. Tolsa and A. Volberg, *The Riesz transform, rectifiability, and removability for Lipschitz harmonic functions*, Preprint, 2012, arXiv:1212.5431, 15 p. To appear in Publ. Mat.

[Po] Ch. Pommerenke, Boundary behaviour of conformal mappings. Springer, Grundlehren der mathematischen Wissenschaften Volume 299, 1992.

[To1] X. Tolsa, *Painlevé's problem and the semiadditivity of analytic capacity*, Acta Math. **190** (2003), no.1, 105–149.

[To2] X. Tolsa, *Painlevé's problem and analytic capacity*, Proceedings of the 7th International Conference on Harmonic Analysis and Partial Differential Equations El Escorial, Madrid (Spain), June 21–25, 2004, Collect. Math. (2006), 89–125.

[To3] X. Tolsa, *Bilipschitz maps, analytic capacity, and the Cauchy integral.* Ann. of Math. (2) 162 (2005), no. 3, 1243–1304.

[To] X. Tolsa, *Analytic capacity, the Cauchy transform, and non-homogeneous Calderón-Zygmund theory.* Progr. Math., vol. **307**, Birkhäuser/Springer, New York 2014, 390 pp.; http://www.mat.uab.cat/∼ xtolsa/llibreweb.pdf.

[Vo] A. Volberg, *Calderón-Zygmund capacities and operators on nonhomogeneous spaces.* CBMS Regional Conf. Ser. in Math. 100, Amer. Math. Soc., Providence, 2003.

[VE] A. Volberg, V. Eiderman, *Non-homogeneous harmonic analysis: 16 years of development*, Uspekhi Mat. Nauk (Russian Math. Surveys), **68** (2013), no. 6 (414), 3–58.

Vladimir Eiderman, Indiana University, Bloomington, Indiana, USA
E-mail address: veiderma@indiana.edu

Alexander Reznikov, Michigan State University, East Lansing, Michigan, USA
E-mail address: rezniko2@msu.edu

Alexander Volberg, Michigan State University, East Lansing, Michigan, USA
E-mail address: volberg@math.msu.edu